

Lab Results

Printed On Jan 14, 2011

Patients with pituitary disease and neuropsychiatric disorders may require endocrinologic consultations because TSH measurements may be misleading and additional testing is needed.

Patients taking the drug AMIODARONE (Cordarone) sometimes have physiologically altered/atypical thyroid test values. Please consult a PDR for a detailed discussion of this.

----- TOXICOLOGY I (URINE) -----

URINE	METHADO	MORPHIN	CODEINE	BARBS	AMPHET	OPIATES
Ref range low	Low Cutoff	NEG		Neg Cutoff	Neg Cutoff	Neg Cutoff
Ref range high	<300 ng/mL	<300 ng/mL		<300 ng/mL	<1000 ng/mL	<300 ng/mL

a 04/19/2010 09:40				neg	neg	neg
b 02/26/2009 15:30				neg	neg	neg
c 02/17/2009 15:38				neg	neg	neg
d 02/10/2009 16:59				neg	neg	neg
e 09/09/2008 05:45				NEGATIVE	NEGATIVE	NEGATIVE
f 12/11/2006 15:35				NEGATIVE	NEGATIVE	NEGATIVE
g 06/26/2006 15:45				NEGATIVE	NEGATIVE	NEGATIVE
h 11/30/1993 15:30	neg	neg	neg	neg		

URINE	COCAINE	MARIJUA	PCP	PHENOTH	PROPOXY
Ref range low	Neg Cutoff	Neg Cutoff	Neg Cutoff		
Ref range high	<300 ng/mL	<100 ng/mL	<25 ng/mL		

a 04/19/2010 09:40	neg	neg			
b 02/26/2009 15:30	neg	neg			
c 02/17/2009 15:38	neg	neg			
d 02/10/2009 16:59	neg	neg			
e 09/09/2008 05:45	NEGATIVE	NEGATIVE	NEGATIVE		
f 12/11/2006 15:35	NEGATIVE	NEGATIVE	NEGATIVE		
g 06/26/2006 15:45	NEGATIVE	NEGATIVE	NEGATIVE		
h 11/30/1993 15:30	neg	neg			

a. Evaluation for COCAINE:

Benzoylcegonine will be detectable in urine after 2-3 hours and urine remains positive for 2-3 days (longer for chronic abusers) after cocaine use.

Evaluation for OPIATES:

Opiate methodology not sensitive to oxycodone. If clinically applicable, oxycodone can be ordered separately.

- b. *** For test BARBS Normals: - ***
 *** For test COCAINE Normals: - ***

PATIENT NAME AND ADDRESS (Mechanical imprinting, if available)

CIAMPI, JOSEPH
 PO BOX 1681
 PALO ALTO, CALIFORNIA 94302

VISTA Electronic Medical Documentation

Printed at PALO ALTO HCS